

Review

Open Access

Neuroprotective and immune effects of active forms of vitamin D₃ and docosahexaenoic acid in Alzheimer disease patients

Milan Fiala¹ and Mathew T. Mizwicki¹

¹Department of Medicine, David Geffen School of Medicine at UCLA and VA Greater Los Angeles Healthcare System, 650 Charles E. Young Dr. South, Los Angeles, CA 90095-1735, USA.

Correspondence: Milan Fiala, M.D., 650 Charles E. Young Dr. South, Room 23-338, Los Angeles, CA 90095-3517, USA

Submission date: November 8, 2011; Acceptance date: December 5, 2011; Publication date: December, 30, 2011

ABSTRACT:

Neurodegenerative diseases, in particular Alzheimer disease (AD), afflict an increasing proportion of the older population with aging. Decreased exposure to sunlight and decreased consumption of fish, fruits, and vegetables, are two epidemiological factors that appear to be related to the pandemic of AD. In addition to replacing simple with complex carbohydrates and avoiding saturated fat, two nutritional components, vitamin D (acting through the endogenous hormonal form 1 α ,25 dihydroxyvitamin D, 1,25D) and docosahexaenoic acid (DHA) (acting through the docosanoid lipidic modulators resolvins and neuroprotectins) have high potential for prevention of Alzheimer disease. 1,25D is a neuroprotective, it acts both directly and indirectly in neurons by improving the clearance of amyloid-beta by macrophages/microglia. Resolvins and neuroprotectins inhibit amyloidogenic processing of amyloid-precursor protein, inflammatory cytokines, and apoptosis. It is likely that the increased consumption of vitamin D and fish oil could prevent neurodegeneration in some subjects by maintaining adequate endocrine, paracrine, and/or autocrine production of 1,25D and the DHA-derived lipidic modulators. Before firm recommendations of the dosage can be proposed, however, the in vivo effects of vitamin D₃ and DHA supplementation should be investigated by prospective studies.

Key words: Alzheimer disease, vitamin D₃, DHA, fish oil

REVIEW:

Alzheimer disease (AD) is the leading cause of dementia worldwide. The major neuropathology of AD is related to amyloid- β 1-42 (A β), a peptide cleaved from the Amyloid precursor protein. The upstream causes of the most common sporadic form of AD are thought to include oxidative stress, dyslipidemia, insulin resistance, physical and psychological stress, and lack of physical

and mental exercise in subjects with the E4 allele of the lipid transport protein apolipoprotein E (ApoE) [1]. The downstream mechanisms of AD are considered to include (a) brain amyloidosis with both fibrillar and oligomeric A β with related synapse loss [2], as described by the amyloid hypothesis [3]; (b) hyperphosphorylation of tau leading to neurofibrillar tangles [4]; and (c) brain inflammation [5]. Although no disease-modifying therapy is available, preventive therapy could address the epidemiological factors associated with AD, i.e. decreased exposure to sunlight, and decreased consumption of fish, fruits, and vegetables.

As recommended for optimal health, nutritional prevention of AD should include a diet with a low amount of saturated fat, and high proportion of complex carbohydrates from cereals and vegetables, such as the Mediterranean diet, with the addition of phytochemicals and other functional food components. In fact, greater adherence to the Mediterranean diet and higher physical activity were independently associated with reduced risk of AD [6]. Anti-oxidants, such as vitamin C and vitamin E, should be included in the diet in physiological amounts [7]. A low cholesterol diet is important because the cholesterol oxidation products, i.e. oxysterols, have pro-inflammatory properties [8]. Patients with diabetes mellitus type II have an increased risk of AD, which could be related to a diet high in calories, or to the metabolic complications of diabetes [9].

In principle, phytochemicals could be protective against AD due to their antioxidant, anti-inflammatory and hypolipidemic properties, including inhibition of (a) protein kinases, such as Akt/protein kinase B (Akt/PKB), Janus kinase 1 (JAK1), glycogen synthase kinase 3 (GSK) [10] [11], (b) dysregulated insulin signaling [12], and (c) cytokines stimulating phosphoinositide 3 kinase (PI3K), GSK, and protein kinase C (PKC) [13].

Chronic diseases are complex and regulated by multiple genes mandating careful and multitargeted selection of phytochemicals. Although there was initial excitement about curcuminoids based on epidemiological data in India and Singapore [14], curcuminoids were not effective in a double-blind trial in AD patients at UCLA [15]. It has become clear that molecular and immunological effects of phytochemicals need to be understood before they can be used for therapy. For example, while curcuminoids have beneficial immune effects on A β phagocytosis at 0.1 μ M, they cause apoptosis at 1 μ M [16]. Furthermore, the heterogeneity of AD mandates that therapy should be personalized in patients with macrophages of different types (Type 0, I and II), as discussed below. Nonetheless, because immune cells migrate from the blood into the brain, improvement of immune cell function stands out as a therapeutic target.

Immunopathology and immunotherapy for Alzheimer disease: Brain amyloidosis is a neuropathological hallmark of AD, and clearance of brain amyloidosis has been the goal of therapies by A β vaccine and anti- A β antibodies [17]. The vaccine and antibodies are designed to stimulate the adaptive immune system for the enhancement of A β phagocytosis by microglia and macrophages. The immunopathology of AD patients also involves the innate immune system, which has not been the prime target of the vaccine. The macrophages of normal subjects are able to phagocytize and degrade A β , but the macrophages of AD patients are defective in phagocytosis and degradation, specifically of A β (not of bacteria) [18]. Although the A β vaccine effectively increased A β clearance in animal models, it failed in patients due to pathological

effects related to tau and amyloid angiopathy [19], lack of clinical efficacy, and encephalitis in 6% of patients [20]. Thus, improving the innate immune system of patients is a novel approach to AD.

We have observed that 1,25D₃, DHA and DHA-derived lipid modulators are effective *in vitro* in recovering dysregulated phagocytosis of A β by AD macrophages. However, the benefits of supplementation with vitamin D₃ or DHA *in vivo* have not been uniformly positive in prospective studies, suggesting that the enzymes responsible for the anabolic production of 1,25D₃ and resolvins by PBMCs/macrophages may be defective in the patients with neurodegenerative diseases. In this respect, we have observed a significant down-regulation of the enzyme 1 α -OHase in PBMCs of a patient with amyotrophic lateral sclerosis (treated with SOD-1), when compared to controls treated in an identical manner. Therefore, nutritional therapy may need to be personalized.

Neuroprotective and immune effects of vitamin D and curcuminoids against dementia:

Numerous studies have shown an association of vitamin D insufficiency with dementia and cognitive decline [21]. Causal relation to dementia was suggested in two prospective studies: the study of 858 older Italian adults indicated relative risk (RR) of cognitive decline (RR 1.6) in subjects with severe 25 (OH)vitamin D insufficiency (<25nM) [22]; the study of 1136 older US men found increased odds of cognitive decline (OR 1.41) in those with \leq 49 nM 25 (OH)vitaminD [23].

1,25D has several intra-cranial mechanisms important for brain health. 1,25D modulates, through the nuclear vitamin D receptor (VDR), the transcription of as many as 500-1000 genes [24]. 1,25D₃ protects against neurotoxicity of A β by downregulating in cortical neurons through VDR L-type voltage-sensitive calcium channels, LVSCC-A1C and LVSCC-A1D, and upregulating nerve growth factor (NGF) and VDR [25]. The role of VDR suppression in AD was highlighted by the demonstration that VDR suppression by siRNA results in an increase of LVSCC-A1C and a decrease of NGF production without a change in LVSCCA1D [26].

As mentioned, the extra-cranial mechanisms of 1,25D₃ involve, in particular, the innate immune system cells, macrophages, and microglia. Macrophages are immune cells that locally produce 1,25D, given that they, like keratinocytes, contain all components of the vitamin D phytoendocrine system [27]. 1,25D regulates anti-microbial macrophage function through potentiation of interferon-gamma production by T cells [28]. 1,25D₃ repairs the deficiency in phagocytosis of A β by AD macrophages through VDR genomic as well as nongenomic mechanisms [16]. *MGAT3 (GnT3)* is an essential gene for phagocytosis of A β by macrophages, as demonstrated using *MGAT3 (GnT3)* siRNA to block A β phagocytosis. We have found that the macrophages of AD patients could be distinguished according to *MGAT3 (GnT3)* transcriptional responses to A β into Type I, II and 0 as follows: Type I macrophages down regulate *MGAT3* and *TLR-3* but are stimulated by curcuminoids to up regulate these genes; Type II macrophages up regulate *MGAT3* and *TLR-3*, but these genes are down regulated by curcuminoids; Type 0 macrophages are transcriptionally –inhibited and do not respond to stimulation [29]. 1,25D₃ increased phagocytosis of A β by macrophages of both Type I and Type II patients. Bisdemethoxycurcumin had small additive effect only in Type I cells [16].

The *in vitro* mechanisms of 1,25D₃ involve potentiation of the currents of the chloride channel named CIC-3 in both Type I and II macrophages. Furthermore, 1,25D₃ elicits a stronger up-regulation of CIC-3 and VDR mRNA's in Type II cells than Type I cells, and possibly has a stronger effect on A β phagocytosis (Mizwicki, Fiala et al, JAD in press). Although there is no firm data on the differences between patients with Type I and II macrophages *in vivo*, anecdotal observations suggest that patients differ in responses to vitamin D₃ supplementation [29].

Regarding the dosage and adverse effects of vitamin D, hypercalcemia has been of concern, however, this has not been documented with administration of vitamin D₃ < 10,000 IU/day (1 IU= 25 ng vitamin D₃) or whole body sunlight exposure (equivalent to 10,000 -20,000 IU vitamin D₃). The dose of vitamin D₃ recommended by the Institute of Medicine is 600 IU before the age 70 years and 800 IU after the age 70 years. To the subjects at risk of deficiency because of race, pregnancy, disease states, obesity, medications, or old age, a higher dose of vitamin D₃ should be administered [30]. In these subjects, at least 1,500 - 2000 IU per day are necessary to maintain vitamin D sufficiency (25-hydroxyvitamin D₃ serum level \geq 30 ng/ml), but higher doses may be beneficial. In our experience, an AD patient with Type II macrophages has been relatively cognitively stable (Mini-Mental State Examination score (MMSE), 19-20) for 3 years on daily supplementation with 5,000 I.U. of vitamin D₃ and 1 gm DHA [29].

Neuroprotective mechanisms of docosahexaenoic acid, resolvin D1 and neuroprotectin D1:

Docosahexaenoic acid (DHA; 4Z,7Z,10Z,13Z,16Z,19Z-docosa-4,7,10,13,16,19-hexaenoic acid) 22:6(n-3) is a major omega-3 fatty acid in neurons, which are essential for brain growth and development. In the brain, DHA has effects on membrane fluidity, synaptic function, and structural integrity and has anti-inflammatory activity [31] via inhibition of calcium channels excitotoxicity [32], and activation of selective gene transcription [33]. In an observational study, high intake of DHA increased gray matter in corticolimbic circuitry [34]. DHA is a precursor for the anti-inflammatory lipid mediators called resolvins, protectins, and neuroprotectins, which have recently been reviewed [35-37]. The E series resolvins, e.g. resolvin E1 (RvE1; 5S,12R,18R-trihydroxy-6Z, 8E, 10E, 14Z, 16E eicosapentaenoic acid) are biosynthesized from the omega-3 fatty acid eicosapentaenoic acid (EPA) 20:5(n-3), and the D series resolvins, e.g. resolvin D1 (RvD1; 7S,8R,17S-trihydroxy-4Z,9E,11E,13Z,15E,19Z-docosahexaenoic acid) from DHA by transcellular synthesis facilitated by aspirin in vascular, leukocytic, and neural cells [38]. Neuroprotectin D1 (NpD1; 10R,17S-dihydroxy-docosa-4Z,7Z,11E,15E,19Z hexaenoic acid) is a potent lipidic mediator induced in the brain after injury, oxidative stress, or stimulation by neurotrophins, which release DHA through phospholipase A2 (PLA2) from neuronal membrane phospholipids. NPD1 is generated from DHA by 15-lipoxygenase-1, epoxidation and hydrolysis. NPD1 has multiple neuroprotective activities, such as down regulation of inflammatory genes, attenuation of leukocyte infiltration, inhibition of apoptosis, and neurotrophic effects through amyloid- β precursor protein (APP)-derived peptide called sAPP α via a non-amyloidogenic, α -secretase-mediated pathway [39] [40]. On the other hand, arachidonic acid (omega-6) in neuronal membranes generates through cyclooxygenase-2 (COX-2) the pro-inflammatory mediators prostaglandin E₂ (PGE₂), prostaglandin I₂ (PGI₂), and thromboxane A₂ [41]. The binding of PGE₂ to the E1 receptor is neurotoxic through disruption of

calcium homeostasis, whereas binding of PGE₂ to the EP2 receptor is neuroprotective [42]. Other mechanisms of DHA include reducing A β production by increasing transthyretin and suppression of phosphorylation of tau [43].

DHA may alter amyloidogenic processing by facilitating the interaction of α -secretase with APP and processing of APP through the nonamyloidogenic pathway [44]. DHA activates pro-survival signaling of Akt by membrane translocation [45]. DHA interacts with the nuclear receptors called retinoid X receptors (RXRs) and peroxisome proliferator-activated receptor- γ (PPAR- γ). RXR/PPAR- γ heterodimers inhibit multiple inflammatory mediators [46].

In some clinical studies, DHA has been shown to be neuroprotective, including traumatic brain injury [47], stroke [48], and cognitive decline in 14 of 17 observational studies [49]. Supplementation with omega-3 fatty acids for 6 months reduced release of IL-1 β and IL-6 from PBMC's of AD patients [50], but this was refuted in a subsequent study [51]. In randomized, double blind clinical trials the overall results of DHA supplementation vs. placebo have been negative, but suggestive of efficacy in early cases and especially in Apo ϵ 4 non-carriers. However, in a recent 18-month prospective study algal DHA supplementation, however without attention to n-6/n-3 ratio, did not slow the rate of cognitive and functional decline in patients with mild to moderate AD (MMSE scores, 14-26), although restricted effect on the ADAS-cog and MMSE was observed in Apo ϵ 4 non-carriers [52]. Thus, the confounding factors in the studies of omega-3 fatty acids are believed to be related to: (a) the degree of neurodegeneration, (b) high dietary n-6 intake, (c) Apo E4 status, (d) oxidation status of the omega-3 fatty acids [49]. The source and stability of the omega 3 fatty acids may be of importance. The company Smartfish (Oslo, Norway) have tested their drink composition which contains omega 3 from fresh locally harvested fish, and found no oxidation products for as long as 1 year after expiration.

Nutritional supplementation with DHA has had greatest impact on cardiovascular prevention, especially prevention of arrhythmias, when the n-6/n-3 ratio was lowered to 1 [41]. The dietary recommendations for lowering the ratio include eliminating N-6 (linoleic acid)-rich foods (e.g. corn, peanut, soybean, sunflower and safflower oils, margarine, and meats), and increasing N-3 (alpha-linolenic)-rich foods (canola oil, linseed oil) and especially N-3 and EPA+DHA-rich foods, (herring, salmon, and tuna with low mercury content). There have been caveats of high serum DHA level association with increased risk of high-grade prostate cancer [53] and with dementia [54]. However, DHA serum levels do not determine the health consequences, because the derivation of pro- and anti-inflammatory mediators from DHA depends on local enzymes in tissues. Nevertheless, administration of polyunsaturated fatty acids, in particular omega-6 but also omega-3, is not without problems: non-enzymatic oxidation of EPA and DHA leads to the formation of isoprostanes and further chain cleavage yields aldehydes, such as malondialdehyde (MDA), acrolein, and hydroxyalkenals, such as 4-hydroxynonenal, which can impair the function of nervous cells [8].

Clinical use of vitamin D₃ and DHA: Although the effects in animal models provide perspective on biochemical mechanisms, such mechanisms have to be shown to be relevant to human patients. As mentioned, AD patients have defects in the innate immune system for A β phagocytosis, which have not been reproduced in animal models. In addition a mouse VDR

responds differently to VDR transcriptional modulators than does the human VDR [55]. For example the MK analogue that blocks $1,25D_3$ transactivation in a number of different cell types and $1,25D_3$ -induced phagocytosis of $A\beta$ by AD macrophages, does not block $1,25D_3$ -VDR transactivation in rodent cells [56]. Furthermore, it has been demonstrated that regulation of the cathelicidin antimicrobial peptide (CAMP) gene by VDR and its ligand $1,25D_3$ is not evolutionarily conserved in mice, rats, or dogs, because the promoters of their CAMP genes lack a VDR response element (VDRE) [57]. Therefore, it is quite clear that the results in mouse model cannot be applied to AD patients and must be determined in prospective studies of persons at risk.

CONCLUSIONS:

The combined use of vitamin D_3 and DHA is an emerging novel strategy to enhance direct and immune protection of neurons against brain amyloidosis and other brain insults. These two compounds will likely have preventive effects in the early stage of neurodegeneration, rather than therapeutic effects.

Competing interests:

Milan Fiala, M.D: Meeting support by Smartfish, Inc.

Authors' contributions:

Both authors contributed equally to this work

REFERENCES:

1. Mahley RW, Weisgraber KH, Huang Y (2006) Apolipoprotein E4: a causative factor and therapeutic target in neuropathology, including Alzheimer's disease. *Proc Natl Acad Sci U S A* 103, 5644-5651.
2. Masliah E, Mallory M, Alford M, DeTeresa R, Hansen LA, McKeel DW, Jr., Morris JC (2001) Altered expression of synaptic proteins occurs early during progression of Alzheimer's disease. *Neurology* 56, 127-129.
3. Hardy J, Selkoe DJ (2002) The amyloid hypothesis of Alzheimer's disease: progress and problems on the road to therapeutics. *Science* 297, 353-356.
4. Gong CX, Grundke-Iqbal I, Iqbal K (2010) Targeting tau protein in Alzheimer's disease. *Drugs Aging* 27, 351-365.
5. Akiyama H, Barger S, Barnum S, Bradt B, Bauer J, Cole GM, Cooper NR, Eikelenboom P, Emmerling M, Fiebich BL, Finch CE, Frautschy S, Griffin WS, Hampel H, Hull M, Landreth G, Lue L, Mrazek R, Mackenzie IR, McGeer PL, O'Banion MK, Pachter J, Pasinetti G, Plata-Salman C, Rogers J, Rydel R, Shen Y, Streit W, Strohmeyer R, Tooyoma I, Van Muiswinkel FL, Veerhuis R, Walker D, Webster S, Wegrzyniak B, Wenk G, Wyss-Coray T (2000) Inflammation and Alzheimer's disease. *Neurobiol Aging* 21, 383-421.
6. Scarmeas N, Luchsinger JA, Schupf N, Brickman AM, Cosentino S, Tang MX, Stern Y (2009) Physical activity, diet, and risk of Alzheimer disease. *JAMA* 302, 627-637.

7. Morris MC, Tangney CC (2011) A potential design flaw of randomized trials of vitamin supplements. *JAMA* 305, 1348-1349.
8. Corsinovi L, Biasi F, Poli G, Leonarduzzi G, Isaia G (2011) Dietary lipids and their oxidized products in Alzheimer's disease. *Mol Nutr Food Res* 55 Suppl 2, S161-SS172.
9. Peila R, Rodriguez BL, Launer LJ (2002) Type 2 diabetes, APOE gene, and the risk for dementia and related pathologies: The Honolulu-Asia Aging Study. *Diabetes* 51, 1256-1262.
10. Aggarwal BB, Shishodia S (2006) Molecular targets of dietary agents for prevention and therapy of cancer. *Biochem Pharmacol* 71, 1397-1421.
11. Takada Y, Fang X, Jamaluddin MS, Boyd DD, Aggarwal BB (2004) Genetic deletion of glycogen synthase kinase-3beta abrogates activation of IkappaBalpha kinase, JNK, Akt, and p44/p42 MAPK but potentiates apoptosis induced by tumor necrosis factor. *J Biol Chem* 279, 39541-39554.
12. McGarry JD (2002) Banting lecture 2001: dysregulation of fatty acid metabolism in the etiology of type 2 diabetes. *Diabetes* 51, 7-18.
13. Permana PA, Menge C, Reaven PD (2006) Macrophage-secreted factors induce adipocyte inflammation and insulin resistance. *Biochem Biophys Res Commun* 341, 507-514.
14. Ng TP, Chiam PC, Lee T, Chua HC, Lim L, Kua EH (2006) Curry consumption and cognitive function in the elderly. *Am J Epidemiol* 164, 898-906.
15. Ringman JM, Frautschy SA, Cole GM, Masterman DL, Cummings JL (2005) A potential role of the curry spice curcumin in Alzheimer's disease. *Curr Alzheimer Res* 2, 131-136.
16. Masoumi A, Goldenson B, Ghirmai S, Avagyan H, Zaghi J, Abel K, Zheng X, Espinosa-Jeffrey A, Mahanian M, Liu PT, Hewison M, Mizwicki M, Cashman J, Fiala M (2009) 1alpha,25-dihydroxyvitamin D₃ Interacts with Curcuminoids to Stimulate Amyloid-beta Clearance by Macrophages of Alzheimer's Disease Patients. *J Alzheimers Dis* 17, 703-717.
17. Galimberti D, Scarpini E (2011) Disease-modifying treatments for Alzheimer's disease. *Ther Adv Neurol Disord* 4, 203-216.
18. Fiala M, Liu PT, Espinosa-Jeffrey A, Rosenthal MJ, Bernard G, Ringman JM, Sayre J, Zhang L, Zaghi J, Dejbakhsh S, Chiang B, Hui J, Mahanian M, Baghaee A, Hong P, Cashman J (2007) Innate immunity and transcription of MGAT-III and Toll-like receptors in Alzheimer's disease patients are improved by bisdemethoxycurcumin. *Proc Natl Acad Sci U S A* 104, 12849-12854.
19. Nicoll JA, Barton E, Boche D, Neal JW, Ferrer I, Thompson P, Vlachouli C, Wilkinson D, Bayer A, Games D, Seubert P, Schenk D, Holmes C (2006) Abeta species removal after abeta42 immunization. *J Neuropathol Exp Neurol* 65, 1040-1048.
20. Maarouf CL, Daugs ID, Kokjohn TA, Kalback WM, Patton RL, Luehrs DC, Masliah E, Nicoll JA, Sabbagh MN, Beach TG, Castano EM, Roher AE (2010) The biochemical aftermath of anti-amyloid immunotherapy. *Mol Neurodegener* 5, 39.
21. Dickens AP, Lang IA, Langa KM, Kos K, Llewellyn DJ (2011) Vitamin D, cognitive dysfunction and dementia in older adults. *CNS Drugs* 25, 629-639.
22. Llewellyn DJ, Lang IA, Langa KM, Muniz-Terrera G, Phillips CL, Cherubini A, Ferrucci L, Melzer D (2010) Vitamin D and risk of cognitive decline in elderly persons. *Arch Intern Med* 170, 1135-1141.

23. Slinin Y, Paudel ML, Taylor BC, Fink HA, Ishani A, Canales MT, Yaffe K, Barrett-Connor E, Orwoll ES, Shikany JM, Leblanc ES, Cauley JA, Ensrud KE (2010) 25-Hydroxyvitamin D levels and cognitive performance and decline in elderly men. *Neurology* 74, 33-41.
24. Haussler MR, Haussler CA, Whitfield GK, Hsieh JC, Thompson PD, Barthel TK, Bartik L, Egan JB, Wu Y, Kubicek JL, Lowmiller CL, Moffet EW, Forster RE, Jurutka PW (2010) The nuclear vitamin D receptor controls the expression of genes encoding factors which feed the "Fountain of Youth" to mediate healthful aging. *J Steroid Biochem Mol Biol* 121, 88-97.
25. Dursun E, Gezen-Ak D, Yilmazer S (2010) A novel perspective for Alzheimer's disease: vitamin D receptor suppression by Amyloid-beta and preventing the Amyloid-beta induced alterations by vitamin D in cortical neurons. *J Alzheimers Dis*.
26. Gezen-Ak D, Dursun E, Yilmazer S (2011) The effects of vitamin D receptor silencing on the expression of LVSCC-A1C and LVSCC-A1D and the release of NGF in cortical neurons. *PLoS One* 6, e17553.
27. Bikle D (2008) Nonclassic Actions of Vitamin D. *J Clin Endocrinol Metab*.
28. Fabri M, Stenger S, Shin DM, Yuk JM, Liu PT, Realegeno S, Lee HM, Krutzik SR, Schenk M, Sieling PA, Teles R, Montoya D, Iyer SS, Bruns H, Lewinsohn DM, Hollis BW, Hewison M, Adams JS, Steinmeyer A, Zugel U, Cheng G, Jo EK, Bloom BR, Modlin RL (2011) Vitamin D Is Required for IFN- γ -Mediated Antimicrobial Activity of Human Macrophages. *Sci Transl Med* 3, 104ra102.
29. Fiala M, Mahanian M, Rosenthal M, Mizwicki MT, Tse E, Cho T, Sayre J, Weitzman R, Porter V (2011) MGAT3 mRNA: A Biomarker for Prognosis and Therapy of Alzheimer's Disease by Vitamin D and Curcuminoids. *J Alzheimers Dis* 25, 135-144.
30. Holick MF, Binkley NC, Bischoff-Ferrari HA, Gordon CM, Hanley DA, Heaney RP, Murad MH, Weaver CM (2011) Evaluation, treatment, and prevention of vitamin D deficiency: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab* 96, 1911-1930.
31. Calder PC (2006) n-3 polyunsaturated fatty acids, inflammation, and inflammatory diseases. *Am J Clin Nutr* 83, 1505S-1519S.
32. Danthi SJ, Enyeart JA, Enyeart JJ (2005) Modulation of native T-type calcium channels by omega-3 fatty acids. *Biochem Biophys Res Commun* 327, 485-493.
33. Jump DB (2008) N-3 polyunsaturated fatty acid regulation of hepatic gene transcription. *Curr Opin Lipidol* 19, 242-247.
34. Conklin SM, Gianaros PJ, Brown SM, Yao JK, Hariri AR, Manuck SB, Muldoon MF (2007) Long-chain omega-3 fatty acid intake is associated positively with corticolimbic gray matter volume in healthy adults. *Neurosci Lett* 421, 209-212.
35. Bannenberg G, Serhan CN (2010) Specialized pro-resolving lipid mediators in the inflammatory response: An update. *Biochim Biophys Acta* 1801, 1260-1273.
36. Serhan CN, Krishnamoorthy S, Recchiuti A, Chiang N (2011) Novel Anti-Inflammatory - Pro-Resolving Mediators and Their Receptors. *Curr Top Med Chem*.
37. Bazan NG, Molina MF, Gordon WC (2011) Docosahexaenoic Acid signalolipidomics in nutrition: significance in aging, neuroinflammation, macular degeneration, Alzheimer's, and other neurodegenerative diseases. *Annu Rev Nutr* 31, 321-351.

38. Serhan CN, Hong S, Gronert K, Colgan SP, Devchand PR, Mirick G, Moussignac RL (2002) Resolvins: a family of bioactive products of omega-3 fatty acid transformation circuits initiated by aspirin treatment that counter proinflammation signals. *J Exp Med* 196, 1025-1037.
39. Lukiw WJ, Cui JG, Marcheselli VL, Bodker M, Botkjaer A, Gotlinger K, Serhan CN, Bazan NG (2005) A role for docosahexaenoic acid-derived neuroprotectin D1 in neural cell survival and Alzheimer disease. *J Clin Invest* 115, 2774-2783.
40. Palacios-Pelaez R, Lukiw WJ, Bazan NG (2010) Omega-3 essential fatty acids modulate initiation and progression of neurodegenerative disease. *Mol Neurobiol* 41, 367-374.
41. Russo GL (2009) Dietary n-6 and n-3 polyunsaturated fatty acids: from biochemistry to clinical implications in cardiovascular prevention. *Biochem Pharmacol* 77, 937-946.
42. Kawano T, Anrather J, Zhou P, Park L, Wang G, Frys KA, Kunz A, Cho S, Orio M, Iadecola C (2006) Prostaglandin E2 EP1 receptors: downstream effectors of COX-2 neurotoxicity. *Nat Med* 12, 225-229.
43. Cole GM, Ma QL, Frautschy SA (2010) Dietary fatty acids and the aging brain. *Nutr Rev* 68 Suppl 2, S102-111.
44. Lukiw WJ, Bazan NG (2010) Inflammatory, apoptotic, and survival gene signaling in Alzheimer's disease. A review on the bioactivity of neuroprotectin D1 and apoptosis. *Mol Neurobiol* 42, 10-16.
45. Akbar M, Calderon F, Wen Z, Kim HY (2005) Docosahexaenoic acid: a positive modulator of Akt signaling in neuronal survival. *Proc Natl Acad Sci U S A* 102, 10858-10863.
46. Tontonoz P, Spiegelman BM (2008) Fat and beyond: the diverse biology of PPARgamma. *Annu Rev Biochem* 77, 289-312.
47. Bailes JE, Mills JD (2010) Docosahexaenoic acid reduces traumatic axonal injury in a rodent head injury model. *J Neurotrauma* 27, 1617-1624.
48. Marcheselli VL, Mukherjee PK, Arita M, Hong S, Antony R, Sheets K, Winkler JW, Petasis NA, Serhan CN, Bazan NG (2010) Neuroprotectin D1/protectin D1 stereoselective and specific binding with human retinal pigment epithelial cells and neutrophils. *Prostaglandins Leukot Essent Fatty Acids* 82, 27-34.
49. Jicha GA, Markesbery WR (2010) Omega-3 fatty acids: potential role in the management of early Alzheimer's disease. *Clin Interv Aging* 5, 45-61.
50. Vedin I, Cederholm T, Freund Levi Y, Basun H, Garlind A, Faxen Irving G, Jonhagen ME, Vessby B, Wahlund LO, Palmblad J (2008) Effects of docosahexaenoic acid-rich n-3 fatty acid supplementation on cytokine release from blood mononuclear leukocytes: the OmegAD study. *Am J Clin Nutr* 87, 1616-1622.
51. Freund-Levi Y, Hjorth E, Lindberg C, Cederholm T, Faxen-Irving G, Vedin I, Palmblad J, Wahlund LO, Schultzberg M, Basun H, Eriksdotter Jonhagen M (2009) Effects of omega-3 fatty acids on inflammatory markers in cerebrospinal fluid and plasma in Alzheimer's disease: the OmegAD study. *Dement Geriatr Cogn Disord* 27, 481-490.
52. Quinn JF, Raman R, Thomas RG, Yurko-Mauro K, Nelson EB, Van Dyck C, Galvin JE, Emond J, Jack CR, Jr., Weiner M, Shinto L, Aisen PS (2010) Docosahexaenoic acid supplementation and cognitive decline in Alzheimer disease: a randomized trial. *JAMA* 304, 1903-1911.

53. Brasky TM, Till C, White E, Neuhouser ML, Song X, Goodman P, Thompson IM, King IB, Albanes D, Kristal AR (2011) Serum phospholipid fatty acids and prostate cancer risk: results from the prostate cancer prevention trial. *Am J Epidemiol* 173, 1429-1439.
54. Laurin D, Verreault R, Lindsay J, Dewailly E, Holub BJ (2003) Omega-3 fatty acids and risk of cognitive impairment and dementia. *J Alzheimers Dis* 5, 315-322.
55. Mizwicki MT, Bula CM, Mahinthichaichan P, Henry HL, Ishizuka S, Norman AW (2009) On the mechanism underlying (23S)-25-dehydro-1alpha(OH)-vitamin D3-26,23-lactone antagonism of hVDRwt gene activation and its switch to a superagonist. *J Biol Chem* 284, 36292-36301.
56. Ishizuka S, Norman AW (1987) Metabolic pathways from 1 alpha,25-dihydroxyvitamin D3 to 1 alpha,25-dihydroxyvitamin D3-26,23-lactone. Stereo-retained and stereo-selective lactonization. *J Biol Chem* 262, 7165-7170.
57. Gombart AF, Borregaard N, Koeffler HP (2005) Human cathelicidin antimicrobial peptide (CAMP) gene is a direct target of the vitamin D receptor and is strongly up-regulated in myeloid cells by 1,25-dihydroxyvitamin D3. *FASEB J* 19, 1067-1077.